

## NEW RECORDS OF THE ISOPOD *Leidya distorta* INFESTING THE MANGROVE CRAB *Ucides cordatus* FROM SOUTH AMERICA\*

Alison Carlos WUNDERLICH<sup>1</sup>, Carlos Alberto CARMONA-SUÁREZ<sup>2</sup>, Marcelo Antonio Amaro PINHEIRO<sup>3</sup>

### ABSTRACT

The Neotropical region has a large diversity of crabs. However, the diversity of the parasites infesting these crustaceans is still poorly known. In this study, we recorded three new occurrences of the isopod bopyrid *Leidya distorta* on *Ucides cordatus* from Southern Brazil and North-Eastern Venezuela. Crabs were collected in mangroves of Venezuela (Cumaná) and Brazil (Cananéia and São Vicente). A total of 281 crabs were captured in these localities and dissected in the laboratory to verify parasitism by *L. distorta*. Three of these crabs were infested with *L. distorta*. Parasitic prevalences in the crabs were generally very low (1.1%), and differed according to localities: Cananéia (2.6%) and São Vicente (2.2%) were slightly higher than in Cumaná (0.5%). Here, we present new records of *L. distorta* infesting the mangrove crab *U. cordatus* from South America and reviewed the brachyuran that serve as host to *L. distorta*.

**Key words:** crab; Brazil; Venezuela; parasitism; bopyridae; isopoda.

## NOVOS REGISTROS DE *Leidya distorta* INFESTANDO O CARANGUEJO-UCÁ *Ucides cordatus* NA AMÉRICA DO SUL

### RESUMO

A região Neotropical tem uma grande diversidade de caranguejos. No entanto, a diversidade de parasitos infestando esses crustáceos ainda é pobremente conhecida. No presente estudo, foram registrados três novas ocorrências de *Leidya distorta* em *Ucides cordatus* no Sudeste do Brasil e no Nordeste da Venezuela. Os caranguejos foram coletados em manguezais da Venezuela (Cumaná) e do Brasil (Cananéia e São Vicente). Um total de 281 caranguejos foram capturados nessas localidades e dissecados no laboratório para verificar a presença de *L. distorta*. Apenas três caranguejos foram infestados com o bopirídeo *Leidya distorta*. A prevalência nos caranguejos foram muito baixa (1.1%), e diferiram de acordo com as localidades: levemente alta em Cananéia (2.6%) e São Vicente (2.2%), comparada com Cumaná (0.5%). Este estudo apresenta novas ocorrências de *Leidya distorta* infestando o caranguejo-uçá *Ucides cordatus* na América do Sul, bem como uma revisão dos braquiúros que servem de hospedeiro para *Leidya distorta*.

**Palavras-chave:** caranguejo; Brasil; Venezuela; parasitismo; bopyridae; isopoda.

**Nota Científica:** Recebido em 24/08/2016; Aprovado em 14/02/2017

<sup>1</sup>UNESP – Universidade Estadual Paulista, Departamento de Parasitologia, Instituto de Biociências de Botucatu (IBB) - Distrito de Rubião Jr., s/n – 18618-970 – Botucatu (SP) – Brasil. E-mail: awunderlich@gmail.com (corresponding author)

<sup>2</sup>Centro de Ecologia, Instituto Venezolano de Investigaciones Científicas, apartado 21827, 1020A, Caracas – Venezuela;

<sup>3</sup>UNESP – Universidade Estadual Paulista, Instituto de Biociências (IB), Campus do Litoral Paulista (CLP), Grupo de Pesquisa em Biologia de Crustáceos (CRUSTA) – Praça Infante Dom Henrique, s/n. – Parque Bitaru – 11330-900 – São Vicente (SP) – Brasil;

\*Fundação de Amparo e Pesquisa do Estado de São Paulo (FAPESP); Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); Instituto Venezolano de Investigaciones Científicas (IVIC).

## INTRODUCTION

Isopods associated with crustacean hosts show a broad distribution around the world (SMIT *et al.*, 2014). According to WILSON (2008), isopods are a diverse group of crustaceans, with more than 10,300 species found in several environments, with 7.7% represented by parasitic isopod epicarideans. Specifically, Bopyridae is represented by 369 genres and 638 species described (WILLIAMS and BOYKO, 2012), being approximately 586 species of this family associated with decapods (BOYKO and WILLIAMS, 2009).

Bopyrid isopods are macroparasites with an indirect life cycle that involves two hosts, both crustaceans (LESTER, 2005). Most of them are found in the branchial chambers of shrimps, for instance, *Probopyrus* spp., (MASUNARI *et al.*, 2000 and GOPALAKRISHNAN *et al.*, 2017) or attached to the pleon of hermit and brachyuran crabs, for example, Athelginae (AN, 2009; WILLIAMS and BOYKO, 2012). According to LESTER (2005), the cryptoniscus that is a free-swimming larvae that seek out a definitive host settles between the gill lamellae

and after moulting a few times, migrates to the roof of the branchial chamber and later mature into a female in a decapod host.

Specifically, the mangrove crab *Ucides cordatus* (Linnaeus, 1763) has been found infested with *Leidyia distorta* (Leidy, 1855) for the first time in the Northeast Brazil (LEMOS DE CASTRO, 1973). *Leidyia distorta*, an isopod species of the family Bopyridae (ROCCATAGLIATA and JORDÁ, 2002; BOYKO *et al.*, 2016) has also been found in crabs of the genus *Uca* spp., from New Jersey, USA (LEIDY, 1855; ROCCATAGLIATA and JORDÁ 2002) to the Rio de La Plata estuary, Argentina (ROCCATAGLIATA and JORDÁ, 2002; JORDÁ and ROCCATAGLIATA, 2002) in Table 1.

However, there is little information about this bopyrid infesting *U. cordatus* in the east coast of South America.

In this study, we recorded three new localities of *L. distorta* infestation in the mangrove crab *U. cordatus*, in two latitudinal separated regions on the Atlantic coast of South America (North-Eastern Venezuela and Southern Brazil), and also reviewed the hosts for this bopyrid species.

**Table 1.** List of brachyuran crabs species used as host by *Leidyia distorta* (Isopoda, Bopyridae).

Hosts	Locality	Region	Reference
<b>Ocypodidae</b>			
<i>Uca pugilator</i>	Atlantic City, New Jersey, USA	NA	Leidy (1855)
	Beaufort, North Carolina, USA	NA	Pearse (1951)
<i>Uca pugnax</i>	Savannah, Georgia, USA	NA	Bourdon & Bowman (1970)
	Guanabara, Rio de Janeiro, Brazil	SA	Lemos de Castro (1973)
<i>Uca minax</i>	Savannah, Georgia, USA	NA	Bourdon & Bowman (1970)
<i>Uca</i> sp.	Sapelo Islands, Georgia, USA	NA	Bourdon & Bowman (1970)
<i>Uca</i> sp.	Port-de-France, Guadeloupe	CA	Bourdon & Bowman (1970)
<i>Uca spinicarpa</i>	Chelem, Yucatan, Mexico	NA	Guillen-Hernandez <i>et al</i> (2013)
<i>Uca vocator</i>	El Sodorro, Trinidad	CA	Bourdon & Bowman (1970)
<i>Uca uruguayensis</i>	Río de la Plata, Argentina	SA	Roccatagliata & Jordá (2002)
<b>Ucididae</b>			
<i>Ucides cordatus</i>	Natal, Rio Grande do Norte, Brazil	SA	Lemos de Castro (1973)
	Caucaia, Ceará, Brazil	SA	Lemos de Castro (1973)
	Cananéia, São Paulo, Brazil	SA	Present study
	São Vicente, São Paulo, Brazil	SA	Present study
	Cumaná, Sucre, Venezuela	SA	Present study

NA, North America; CA, Central America; SA, South America

## METHODS

Mangrove crabs were collected from two localities in Southern Brazil (São Vicente and Cananéia) and one in Venezuela (Cumaná, Sucre State) (Figure 1).


A total of 85 crabs were captured between February 2012 to June 2015 in mangrove areas in both localities (São Vicente: 23°58'51.4"S; 46°24'24.5"W; Cananéia: 24°59'9.80"S; 47°56'12.24"W) in Southern Brazil. The sampling area ranged between 12,500 and 27,500 m<sup>2</sup>, with a predominance of *Rhizophora mangle* L. trees (>70%). In these localities, crabs were caught using "braceamento", a manual method where the fisherman uses his arm to extract the crabs from their burrows (PINHEIRO and FISCARELLI, 2001) and legally permitted by the Brazilian law (BRASIL, 2011).

In Cumaná, the crabs were collected between February and August 2011 at the spillway of the sea close to the Manzanares river, Sucre State, Venezuela (10°27'43.11"N, 64°6'07.99"W). This site had a predominant vegetation of *Avicennia germinans* (L.)

with a scarce presence of *R. mangle* at the border of the spillway. With local fishermen help, fishing net pieces were placed over the burrow entrances. Crabs trying to exit the burrows got entangled in the net and could be easily captured. Afterwards, they were put in an icebox and transported to the laboratory.

In the laboratory, the specimens of *U. cordatus* were sexed by inspection of the abdomen shape and number of pleopods (PINHEIRO and FISCARELLI, 2001), and the carapace width (CW) measured with a precision vernier caliper (0.05 mm). Later, the specimens were sacrificed and searched for bopyrids in their branchial chambers. The collected isopods were identified as *L. distorta* based on the descriptions given by RICHARDSON (1908), BOURDON and BOWMAN (1970), and ROCCATAGLIATA and JORDÁ (2002).

The bopyrids were fixed in 70% ethanol, photographed, and the prevalence was calculated as the number of hosts infested divided by the number of hosts examined expressed as a percentage (BUSH *et al.*, 1997).



**Figure 1.** Map showing three localities from South American region (Cananéia and São Vicente, at São Paulo State, Brazil; and Cumaná, Sucre State, Venezuela) from which *Leidyia distorta* individuals were registered.

## RESULTS

A total of 281 crabs were collected from the three localities (Cananéia, N = 40; São Vicente, N = 45; and Cumaná, N = 196). From this total, three specimens (one of each locality) were infested with different developmental stages of the bopyrid *L. distorta* (Figures 2, 3, and 4). The prevalence of *L. distorta* in *U. cordatus* was very low throughout the study from the three localities, with a mean value of 1.1%. The prevalence was slightly higher for Cananéia (2.6%) and São Vicente (2.2%) when compared to Cumaná (0.5%) (Table 2). One juvenile of *L. distorta* was found in each locality. A total of six male bopyrids and an adult female of *L. distorta* were found only in Cananéia (Table 2), parasitizing a mature adult male of *U. cordatus* (73 mm CW).

The adult female of this species is characterised by possessing a large bilobed head, with a strong dorsal anterior carina (HE, head; Figure 2A). Older mature female possesses a fully expanded and spherical marsupium and well-developed dorsal bosses (Figure 2 A, B). First two pereomere (1-2) are shorter. The three following segments are the largest (3-5) and are sub equal in length, about twice as long as the first. The 6-7<sup>th</sup> pereomeres are a little shorter than the fifth, and the seventh is half as long as the sixth. The seven pairs of pereopods are small and feeble, with the propodus and dactyls forming a prehensile hand. The terminal segment is furnished with a single pair of elongated pinnate appendages, the uropod (Figure 2 C). The pleopods are biramous (Figure 2 C, D), and the antennae are little developed. The maxillipeds with palp are present, articulated or not. There are also five pairs of incubatory plates, which


overlap each other on the ventral side, completely inclosing the marsupial cavity. The pereonites with prominent mid-dorsal projections (i.e., dorsal bosses on the pereon) is an important diagnosis which is characteristic for this species in adult females (Figure 2 A). The juvenile females (Figure 3) differs from the adult in its narrower and elongated form, the thorax not much distended, and more symmetrical segments of the pereon (Figure 3 A), which do not have the mid-dorsal projections as in the adult. They also have the segments of the pereon more distinctly separated and the lateral appendages more elongated (Figure 3 A, B). Also, the oostegites in the juvenile females are much smaller than the ones in the adult females, and the marsupial pouch greatly reduced in size (Figure 3).

The marsupial pouch is never filled with eggs at this stage. The male of this species possessed a body narrow and elongated (Figure 4). Head than long; frontal margins straight, with anterolateral angles rounded (Figure 4 C). The posterior portion is somewhat wider than the anterior (Figure 4 D). Vestiges of ocular pigment visible from posterior part of the head. The first pair of antennae (Figure 4 B) composed of three joints, the second pair composed of seven, the last two being minute and tipped with setae. The seven thoracic segments are separated from each other by lateral incisions (Figure 4 A, B) so that the lateral margins are not continuous. There are seven pairs of prehensile legs. The six abdominal segments are perfectly distinct, separated from each other by lateral incisions, and decrease gradually in width, from the first to the sixth, which carries a pair of elongated, tapering appendages, the uropod (Figure 4 D). The pleopods are rudimentary.


**Table 2.** Infestation levels of *Leidyia distorta* (Isopoda, Bopyridae) registered in three localities from South America in the present study.

Locality	NGC	NPC	Instars of <i>Leidyia distorta</i> *			Prevalence (%)
			Male	Juvenile female	Adult female	
Cananéia, Brazil (MZUSP 35962)**	40	1	6	1	1	2.55
São Vicente, Brazil (MZUSP 35961)**	45	1	-	1	-	2.22
Cumaná, Venezuela	196	1	-	1	-	0.51
Total	281	3	6	3	1	1.07


NGC, numbers of gathered crabs; NPC, numbers of parasitized crabs; \*Number of developmental stages of *Leidyia distorta* according to the description of Roccatagliata and Jordá (2002). \*\*The voucher parasite specimens were deposited in the Museum of Zoology of the University of São Paulo (MZUSP), Brazil.



**Figure 2.** *Leidyia distorta* (Leidy, 1855). Adult female specimen: A, head and well developed dorsal bosses; B, marsupium fully expanded and spherical; C, pleopods; and pleotelson; D, pleopods. Where: DB, dorsal bosses; EM, expanded marsupium; HE, head; PT, pleotelson. Scale bar = 2 mm (A and B), 1 mm (C and D).



**Figure 3.** *Leidyia distorta* (Leidy, 1855). Juvenile female specimen: A, dorsal view; B, ventral view; C, head and lamina view; D, pleopods and pereopods view; and E, antennae and maxilliped view. Where: BP, biramous pleopod; FLT, frontal lamina truncate; HE, head; MDP, mid dorsal projection; MX, maxillipeds; PE, pereopods; UR, uropod. Scale bar = 2 mm (A, B and D), 1 mm (C and E).



**Figure 4.** *Leidyia distorta* (Leidy, 1855). Male specimen: A, dorsal view; B, ventral view; C, head view; D, pair of elongated uropod. Where: AN, antennae; HE, head; PE, pereopods; UR, uropod. Scale bar = 1 mm (A and B), 500 µm (C and D).

## DISCUSSION

In this study, we recorded for the first time the infestation of the mangrove crab *U. cordatus* by *L. distorta* in Venezuela, and it also represents the second report in Brazil. LEMOS DE CASTRO (1973) reported the first record of this isopod infestation previously in *U. cordatus* in two different localities of Northeast Brazil (Natal, Rio Grande do Norte State; and Caucaia, Ceará State). Thus, in the present study, we expanded the distribution of this bopyrid by adding three new localities, two in the Southeast region (Cananéia and São Vicente, São Paulo state, Brazil) and another one in the Northwest of South America (Cumaná, Sucre state, Venezuela).

This bopyrid has also been found in species of the genus *Uca* in several regions of the American continent coast (Table 1), since New Jersey, USA (LEIDY, 1855; ROCCATAGLIATA and JORDÁ, 2002) until Rio de La Plata estuary, Argentina (ROCCATAGLIATA and JORDÁ, 2002; JORDÁ and ROCCATAGLIATA, 2002). From all the known species of the genus *Uca* in Brazil, this bopyrid only infests the fiddler crab *U. pugnax* (Smith) (LEMOS DE CASTRO, 1973). Moreover, according to an important revision about the diversity of parasitic isopods associated with crustacean hosts, the poor

knowledge about Neotropical bopyrids could be due to a lack of sufficient sampling in many of these regions (e.g. east coast of South America), rather than reflecting real biogeographic patterns for bopyrids (WILLIAMS and BOYKO, 2012).

According to LESTER (2005), adult males are much smaller than the females (Figure 4) and are usually found attached to their pleopods. JORDÁ and ROCCATAGLIATA (2002), found a very low prevalence in both sexes of *Uca uruguayensis* Nobili, 1901 (males, 12%; and females, 5%). The data given by these authors are similar, with the lowest prevalence found in the populations of *U. cordatus* in São Vicente (Brazil), Cananéia (Brazil), and Cumaná (Venezuela). LEMOS DE CASTRO (1973), also found a very low prevalence in *U. cordatus* from Rio Grande do Norte and Ceará states, and in *U. pugnax* from Rio de Janeiro, all considered Brazilian sites. Moreover, LAFFERTY and KURIS (2009) point out that the infestation pattern of *L. distorta* is often aggregated, which could explain the lowest prevalence in mangrove crabs. In addition, single infestations in adult female bopyrid isopods are generally accompanied by a single dwarf male consort (ROCCATAGLIATA and JORDÁ, 2002; LAFFERTY and KURIS, 2009). Rarely two parasites can occur together in the branchial chamber or under

the abdomen of the decapod hosts, and never three or more (LAFFERTY and KURIS, 2009), but this fact was not verified in the present study.

Therefore, the low prevalence of bopyrid species in brachyuran crabs and the reported numbers of undescribed species in recent studies suggest a large yet undescribed fauna of marine isopoda, especially in the tropical Atlantic waters (POORE and BRUCE, 2012). Our results revealed a lower prevalence of *L. distorta* infesting the mangrove crab *U. cordatus* from South America.

## ACKNOWLEDGEMENTS

This research was supported in Brazil by FAPESP – “Fundação de Amparo e Pesquisa do Estado de São Paulo” during expeditions of “Projeto Uçá” Phase III (FAPESP, Process # 2009/14725-1) and IV (CNPq), and in Venezuela by IVIC internal funding # 961. We are very thankful to the fishermen, especially to Anacleute Silva (Brazil) and Nicolás Suárez (Venezuela), as well as to Caroline Souza, Nicholas Kriegler (Brazil), Arianny Camejo and Wílmer Rojas (Venezuela) for processing the samples of *U. cordatus*. We also thank Prof. Reinaldo José da Silva (UNESP, IBB) for logistic support. Finally, we thank Flavia P. Zanotto for the English review.

## REFERENCES

- AN, J. 2009 A review of bopyrid isopods infesting crabs from China. *Integrative and Comparative Biology*, 49(2): 95-105.
- BOURDON, R; BOWMAN, T.E. 1970 Western Atlantic species of the parasitic genus *Leidyia* (Epicaridea: Bopyridae). *Proceedings of the Biological Society of Washington*, 83(1): 409-424.
- BOYKO, C.B; BRUCE, N.L.; MERRIN, K. L.; OTA, Y.; POORE, G.C.B.; TAITI, S.; SCHOTTE, M.; WILSON, G.D.F. (Eds) (2008 onwards) 2016 WoRMS Isopoda: World Marine, Freshwater and Terrestrial Isopod Crustaceans database (version 2016-03-01). In: *Species 2000 & ITIS Catalogue of Life, 25th March 2016* (Roskov Y., Abucay L., Orrell T., Nicolson D., Kunze T., Flann C., Bailly N., Kirk P., Bourgoin T., DeWalt R.E., Decock W., De Wever A., eds). Digital

resource at [www.catalogueoflife.org/col](http://www.catalogueoflife.org/col). Species 2000: Naturalis, Leiden, the Netherlands. ISSN 2405-8858.

- BOYKO, C.R.; WILLIAMS, J.D. 2009 Crustacean parasites as phylogenetic indicators in decapod evolution. In: (Eds. J.W. Martin, K.A. Crandall and D.L. Felder) *Crustacean Issues: Decapod Crustacean Phylogenetics*. Taylor and Francis/CRC Press, Boca Raton. p. 197-220.
- BRASIL, 2011 *Proposta do Plano Nacional de Gestão para o Uso Sustentável do Caranguejo-Uçá, do Guaiamum e do Siri-Azul*. Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA), Brasília, Brasil. 156 p.
- BUSH, A.O.; LAFFERTY, K.D.; LOTZ, J.M.; SHOSTAK, A.W. 1997 Parasitology meets ecology on its own terms: Margolis *et al.* Revisited. *Journal of Parasitology*, 83(4): 575-583.
- GOPALAKRISHNAN, A.; RAJA, K.; TRILLES, J.P.; RAJKUMAR, M.; RAHMAN, M.M.; SARAVANAKUMAR, A. 2017 Bopyrid isopods parasitizing on the cultured fresh water prawn, *Macrobrachium malcolmsonii* in South India. *Journal of Parasitic Diseases*, 41(1): 93-96. DOI 10.1007/s12639-016-0756-7.
- GUILLEN-HERNANDEZ, S.; PÉREZ-POVEDANO, M.A.; ZEPEDA-AGUILAR, M.; GONZÁLEZ-SALAS, C.F. 2013 Variación estacional de los helmintos parásitos de *Uca spinicarpa* en tres sitios del sistema Lagunar de Chelem, Yucatán, México. In: Proceedings of the 3rd Symposium sobre el conocimiento de los recursos costeros del Sureste de México: componente biológico, 74, Mérida, México, 27-31 May/2013. Disponível em: < [http://www.conabio.gob.mx/institucion/proyectos/resultados/KU011\\_Anexo\\_Libro\\_Resumenes.pdf](http://www.conabio.gob.mx/institucion/proyectos/resultados/KU011_Anexo_Libro_Resumenes.pdf)> Acesso em: 10 mar. 2015.
- JORDÁ, M.T.; ROCCATAGLIATA, D. 2002 Population dynamics of *Leidyia distorta* (Isopoda: Bopyridae) infesting the fiddler crab *Uca uruguayensis* at Río de La Plata Estuary, Argentina. *Journal of Crustacean Biology*, 22(4): 719-727.
- LAFFERTY, K.D.; KURIS, A.M. 2009 Parasitic

- castration: The evolution and ecology of body snatchers. *Trends in Parasitology*, 25(12): 564–572.
- LEIDY, J. 1855 Contributions toward a knowledge of the marine invertebrate fauna of the coasts of Rhode Island and New Jersey. *Journal of the Academy of Natural Sciences of Philadelphia*, 3(1): 135–152.
- LEMOS DE CASTRO, A. 1973 Crustáceos isópodos epicarídeos do Brasil. VI. *Leidya distorta* (Leidy): distribuição geográfica e hospedeiros. *Boletim do Museu Nacional, Nova Série, Zoologia*, 286(1): 1-5.
- LESTER, R.J.G. 2005 Isopoda (isopods). In: Rohde, K. *Marine Parasitology*. CSIRO Publishing, Wallingford. p. 138-144.
- MASUNARI, S.; CASTAGINI, A.S.; OLIVEIRA, E. 2000 The population structure of *Probopyrus floridensis* (Isopoda, Bopyridae). A parasite of *Macrobrachium potiuna* (Decapoda, Palaemonidae) from the Pereque River Paranagua Basin, South Brazil. *Crustaceana*, 73(9): 1095–1108.
- MCDERMOTT, J.J. 1991 Incidence and host-parasite relationship of *Leidya bimini* (Crustacea, Isopoda, Bopyridae) in the brachyuran crab *Pachygrapsus transversus* from Bermuda. *Ophelia*, 33(2): 71-95.
- PINHEIRO, M.A.A.; FISCARELLI, A.G. 2001 *Manual de apoio à fiscalização do caranguejo-uçá (Ucides cordatus)*. Instituto Brasileiro do Meio Ambiente (IBAMA) / Centro de Pesquisa e Gestão de Recursos Pesqueiros do Litoral Sudeste e Sul (CEPSUL), 1ª Edição, ISBN 85-88570-02-5, Itajaí. 43 p.
- POORE, G.C.B.; BRUCE, N.L. 2012 Global Diversity of Marine Isopods (Except Asellota and Crustacean Symbionts). *PLoS ONE* 7(8): 1-15. doi:10.1371/journal.pone.0043529.
- RICHARDSON, H. 1908 The parasitic isopod *Leidya distorta* (Leidy) found on a new host. *Proceedings of the United States National Museum*, 34(1593): 23-26.
- ROCCATAGLIATA, D.; TORRES-JORDÁ, M. 2002 Infestation of the fiddler crab *Uca uruguayensis* by *Leidya distorta* (Isopoda, Bopyridae) from the Río de la Plata estuary, Argentina. *Journal of Crustacean Biology*, 22(1): 69-82.
- SMIT, N.J.; BRUCE, N.L.; HADFIELD, K.A. 2014 Global diversity of fish parasitic isopod crustaceans of the family Cymothoidae. *International Journal for Parasitology: Parasites and Wildlife*, 3(2): 188-197.
- WILSON, G.D.F. 2008 Global Diversity of isopod crustaceans (Crustacea: Isopoda) in freshwater. In: (Eds. E. V. Balian, C. Lévêque, H. Segers, K. Martens) *Freshwater Animal Diversity Assessment. Hydrobiologia*, 595(1): 231-240.
- WILLIAMS, J.D.; BOYKO, C.B. 2012 The Global Diversity of Parasitic Isopods Associated with Crustacean Hosts (Isopoda: Bopyroidea and Cryptoniscoidea). *PLoS ONE*, 7(4): 1-9.